

Name: Ege Yalcinbas
PI Name: Kristen Maynard

Email: ege.yalcinbas@libd.org
PI Email: kristen.maynard@libd.org

Transcriptional Impact of Chronic Fentanyl Self-Administration on Habenula Circuits

Ege A. Yalcinbas^{1,2}, Daianna Gonzalez-Padilla¹, Emma Chaloux-Pinette⁴, Melissa Mayén-Quiroz¹, Robin Magnard⁵, Beau A. Oster^{1,3}, Leonardo Collado-Torres^{1,6,7}, Patricia H. Janak^{3,5}, and Kristen R. Maynard^{1,2,3}

¹Lieber Institute for Brain Development, Johns Hopkins Medical Campus, Baltimore, MD, 21205, USA; ²Department of Psychiatry and Behavioral Sciences, Johns Hopkins School of Medicine, Baltimore, MD, 21205, USA; ³The Solomon H. Snyder Department of Neuroscience, Johns Hopkins School of Medicine, Baltimore, MD, 21205, USA; ⁴Medical Scientist Training Program, Johns Hopkins University School of Medicine, Baltimore, MD, 21205, USA; ⁵Department of Psychological and Brain Sciences Johns Hopkins University, Baltimore, MD, 21218, USA; ⁶Department of Biostatistics, Johns Hopkins Bloomberg School of Public Health, Baltimore, MD, 21205, USA; ⁷Center for Computational Biology, Johns Hopkins University, Baltimore, MD, 21205, USA

The habenula (Hb) modulates reward processing and pain/aversive states through its connectivity with midbrain structures, notably the dopaminergic ventral tegmental area (VTA). Hb dysfunction has been implicated in a variety of neuropsychiatric disorders, including substance use disorders. Hb expresses high levels of μ -opioid receptors, which mediate the analgesic effects of opioids. However, it is unknown how Hb cell types are transcriptionally impacted in opioid use disorder (OUD). To identify genes regulated by chronic opioid intake, we performed bulk RNA-sequencing (RNA-seq) on Hb tissue from rats exposed to an extended access fentanyl self-administration (SA) paradigm. Differential gene expression analysis identified 453 differentially expressed genes (DEGs, FDR < 0.05) associated with chronic fentanyl SA compared to saline controls. Gene ontology analyses implicated synaptic plasticity, glutamatergic, neuromodulatory, and neuropeptide signaling, and mitochondrial dysfunction. Cell type enrichment analyses using mouse and human single nucleus (sn) RNA-seq data revealed that upregulated DEGs were enriched in a unique lateral Hb neuronal subpopulation. This subpopulation, which was defined by expression of *Kcnmb4*, and also enriched for *Oprm1*, may represent a cell type that is particularly vulnerable to chronic opioid SA. We are now performing cell type- and circuit-specific molecular profiling follow up studies. Molecular genetic labeling of VTA-projecting Hb neurons revealed that a subset expresses *Oprm1*. By combining these circuit labeling techniques with snRNA-seq, we obtained transcriptomic signatures for VTA-projecting Hb neurons and evaluated their enrichment in fentanyl-responsive genes. These studies are expected to provide new molecular insights into the contribution of Hb dysregulation to OUD.