

Submitter Name: Lisa Maturin  
Submitted email: [lmaturin@ucsd.edu](mailto:lmaturin@ucsd.edu)  
PI Name (if different): Olivier George  
PI email (if different): [olgeorge@ucsd.edu](mailto:olgeorge@ucsd.edu)

**The Cocaine and Oxycodone Biobanks: Two repositories of biological samples from genetically characterized outbred rats that exhibit compulsive-like escalation of cocaine or oxycodone self-administration**

Lisa Maturin<sup>1</sup>, Marsida Kallupi<sup>1</sup>, Brent Boomhower<sup>1</sup>; Molly Brennan<sup>1</sup>; Mckenzie Pavlich<sup>1</sup>; Lani Tieu<sup>1</sup>; Abraham Palmer<sup>1</sup>, Leah Solberg Wood<sup>2</sup>, Giordano de Guglielmo<sup>1</sup>, Olivier George<sup>1</sup>

<sup>1</sup>Department of Psychiatry, University of California San Diego, <sup>2</sup>Department of Molecular Medicine Wake Forest University

Identification of the mechanisms that underlie compulsive cocaine or oxycodone use in animal models is a major goal for understanding the genetic risk factors for addiction and facilitating the identification of novel druggable targets. A key issue for the field is the lack of a repository that contains biological samples from behaviorally and genetically characterized rats. We introduce the Cocaine Biobank ([www.cocainebiobank.org](http://www.cocainebiobank.org)) and the Oxycodone Biobank ([www.oxycodonebiobank.org](http://www.oxycodonebiobank.org)), two repositories of biological samples from a unique, genetically diverse strain of outbred heterogeneous stock (HS) rats that have been behaviorally and genetically characterized using next-generation sequencing, state-of-the-art behavioral screening, and a variety of preservation techniques. Male and female rats are trained to self-administer cocaine (0.5 mg/kg/inf) in daily 6 h sessions or oxycodone (0.15 mg/kg/inf) in daily 12 h sessions and tested using progressive-ratio responding, responding despite adverse consequences (contingent footshocks), and measures of analgesia, hyperalgesia and irritability-like behaviors. Results show high individual variability with vulnerable and resistant rats that is likely to facilitate the detection of gene variants and the molecular and cellular mechanisms of addiction. Preservation techniques include perfusion, snap-freezing, and cryopreservation maximize the compatibility of these tissue banks with cellular, molecular, and anatomical methods. The Biobanks provides free access to over 20 organs. The Biobanks have the potential to facilitate identification of novel druggable targets and provide a unique data/tissue repository that will facilitate follow-up and replication studies.